arXiv:1404.5404v1 [math.NT] 22 Apr 2014

Congruences for the number of partitions and bipartitions with distinct even parts

Haobo Dai*

Abstract

Let $ped(n)$ denote the number of partitions of n wherein even parts are distinct (and odd parts are unrestricted). We show infinite families of congruences for $ped(n)$ modulo 8. We also examine the behavior of $ped_{-2}(n)$ modulo 8 in detail where $ped_{-2}(n)$ denotes the number of bipartitions of n with even parts distinct. As a result, we find infinite families of congruences for $ped_2(n)$ modulo 8.

Keywords partitions and bipartitions with even parts distinct; congruences; binary quadratic forms.

Mathematics Subject Classification (2000): 05A17;11P83

1 Introduction

Let $ped(n)$ denote the number of partitions of n wherein even parts are distinct (and odd parts are unrestricted). The generating function for $ped(n)$ ([1]) is

$$\sum_{n=0}^{\infty} ped(n)q^n := \frac{(q^4;q^4)_\infty}{(q;q)_\infty} = \prod_{m=1}^{\infty} \frac{(1-q^{4m})}{(1-q^m)}. \tag{1.1}$$

Note that by (1.1), the number of partitions of n wherein even parts are distinct equals the number of partitions of n with no parts divisible by 4, i.e., the 4-regular

*Department of Mathematics, University of Shanghai Jiao Tong University, Shanghai, 200240, China; e-mail: dedekindbest@hotmail.com

partitions (see [1] and references therein). The arithmetic properties were studied by Andrews, Hirshhorn and Sellers [1] and Chen [3]. For example, in [1], Andrews, et al., proved that for all $n \geq 0$,

$$ped(9n+4) \equiv 0 \pmod 4 \tag{1.2}$$

and

$$ped(9n+7) \equiv 0 \pmod 4. \tag{1.3}$$

Suppose that r is an integer such that $1 \leq r < 8p$, $rp \equiv 1 (\mathrm{mod}\, 8)$, and $(r,p) = 1$. By using modular forms, Chen [3] showed that if $c(p) \equiv 0 (\mathrm{mod}\, 4)$, then, for all $n \geq 0$, $\alpha \geq 1$,

$$ped\left(p^{2\alpha}n + \frac{rp^{2\alpha-1}-1}{8}\right) \equiv 0 \pmod 4 \tag{1.4}$$

where $c(p)$ is the p-th coefficient of $\frac{\eta^4(16z)}{\eta(8z)\eta(32z)} := \sum_{n=1}^{\infty} c(n)q^n$. Note that in [3], Chen didn't show the coefficients of $c(p)$ explicitly. Note also that in a beautiful paper [4], Chen studied arithmetic properties for the number of k-tuple partitions with even parts distinct modulo 2 for any positive integer k by using Hecke nilpotency. Berkovich and Patane [2] calculated $c(n)$ explicitly. In particular, they showed that $c(p) = 0$ if and only if $p = 2$, $p \equiv 5 (\mathrm{mod}\, 8)$ and $p \equiv 3 (\mathrm{mod}\, 4)$.

As a direct application of Chen's, Berkovich and Patane's theorems, we have the following.

Corollary 1.1. *Let p be a prime which is congruent to 5 modulo 8 or congruent to 3 modulo 4 and suppose that r is an integer such that $1 \leq r < 8p$, $rp \equiv 1 (\mathrm{mod}\, 8)$, and $(r,p) = 1$, then*

$$ped\left(p^{2\alpha}n + \frac{rp^{2\alpha-1}-1}{8}\right) \equiv 0 \pmod 4$$

for all $n \geq 0$ and $\alpha \geq 1$.

Ono and Penniston [8] showed an explicit formula for $Q(n)$ modulo 8 by using the arithmetic of the ring of $\mathbb{Z}[\sqrt{-6}]$ where $Q(n)$ denotes the number of partitions of an integer n into distinct parts. We are unable to explicitly determine $ped(n)$ modulo 8. But we can prove infinitely families congruences for $ped(n)$ modulo 8. Our first main result is the following.

Theorem 1.2. *Let p be a prime which is congruent to $7 (\mathrm{mod}\, 8)$. Suppose that r is an integer such that $1 \leq r < 8p$, $rp \equiv 1 (\mathrm{mod}\, 8)$, and $(r, p) = 1$, then for all $n \geq 0$, $\alpha \geq 0$, we have*

$$ped\left(p^{2\alpha+2}n + \frac{rp^{2\alpha-1}+1}{8}\right) \equiv 0 \pmod 8.$$

Example 1.3. For all $n \geq 0$, $\alpha \geq 0$,

$$ped\left(7^{2\alpha+2}n + \frac{r \times 7^{2\alpha+1}-1}{8}\right) \equiv 0 \pmod 8,$$

for $r = 15, 23, 31, 39$ and 47.

Let $ped_{-2}(n)$ be the number of bipartitions of n with even parts distinct. The generating function of $ped_{-2}(n)$ [5] is

$$\sum_{n=0}^{\infty} ped_{-2}(n)q^n := \frac{(q^4;q^4)_\infty^2}{(q;q)_\infty^2} = \prod_{m=1}^{\infty} \frac{(1-q^{4m})^2}{(1-q^m)^2}. \tag{1.5}$$

Recently, in [5], Lin investigated arithmetic properties of $ped_{-2}(n)$. In particular, he showed following theorems:

Theorem 1.4. *([5]) For $\alpha \geq 0$ and any $n \geq 0$, we have*

$$ped_{-2}(n)\left(3^{2\alpha+2}n + \frac{11 \times 3^{2\alpha+1}-1}{4}\right) \equiv 0 \pmod 3,$$

$$ped_{-2}(n)\left(3^{2\alpha+3}n + \frac{5 \times 3^{2\alpha+2}-1}{4}\right) \equiv 0 \pmod 3.$$

Theorem 1.5 ([5])**.** *$ped_{-2}(n)$ is even unless n is of the form $k(k+1)$ for some $k \geq 0$. Furthermore, $ped_{-2}(n)$ is a multiple of 4 if n is not the sum of two triangular numbers.*

As a corollary of Theorem 1.3 and 1.4, Lin proved an infinite family of congruences for $ped_{-2}(n)$ modulo 12:

$$ped_{-2}\left(3^{2\alpha+2}n + \frac{11 \times 3^{2\alpha+1}-1}{4}\right) \equiv 0 \pmod{12},$$

for any integers $\alpha \geq 0$ and $n \geq 0$.

As in [8], our second main achievement is to examine $ped_{-2}(n)$ modulo 8 in detail.

Theorem 1.6. *If n is a non-negative integer, then let N and M be the unique positive integers for which*

$$4n+1 = N^2 M$$

where M is square-free. Then the following are true:

(1) If $M = 1$, then $ped_{-2}(n)$ is odd.

(2) If $M = p$, and $ord_p(4n+1) \equiv 1 (\mathrm{mod}\, 4)$, then $ped_{-2}(n) \equiv 2 (\mathrm{mod}\, 4)$.

(3) If $M = p$, and $ord_p(4n+1) \equiv 3 (\mathrm{mod}\, 8)$, then $ped_{-2}(n) \equiv 4 (\mathrm{mod}\, 8)$.

(4) If $M = p_1 p_2$, where p_1 and p_2 are distinct primes, $p_i \equiv 1 (\mathrm{mod}\, 4)$ and $ord_{p_i}(4n+1) \equiv 1 (\mathrm{mod}\, 4)$ for $i = 1, 2$, then $ped_{-2}(n) \equiv 4 (\mathrm{mod}\, 8)$.

(5) In all other cases we have that $ped_{-2}(n) \equiv 0 (\mathrm{mod}\, 8)$.

As a corollary to Theorem 1.5. we can show infinite families of congruences for $ped_{-2}(n)$ modulo 8.

Corollary 1.7. *(i) For $p \equiv 3 (\mathrm{mod}\, 4)$, and let $1 \leq r < 4p$, $(r, p) = 1$, $rp \equiv 1 (\mathrm{mod}\, 4)$, then we have*

$$ped_{-2}\left(p^{2\alpha+2} n + \frac{r \times p^{2\alpha+1} - 1}{4}\right) \equiv 0 \pmod 8, \tag{1.6}$$

for $\alpha \geq 0$ and any $n \geq 0$.

(ii) For $p \equiv 1 (\mathrm{mod}\, 4)$, and let $1 \leq r < 4p$, $(r, p) = 1$, $rp \equiv 1 (\mathrm{mod}\, 4)$, then we have

$$ped_{-2}\left(p^{8\alpha+8} n + \frac{r \times p^{8\alpha+7} - 1}{4}\right) \equiv 0 \pmod 8, \tag{1.7}$$

for $\alpha \geq 0$ and any $n \geq 0$.

Proof. Since the proof of (i) and (ii) are similar, we only give the proof of (ii). Note that

$$4\left(p^{8\alpha+8} n + \frac{r \times p^{8\alpha+7} - 1}{4}\right) + 1 = p^{8\alpha+7}(4pn + r),$$

so for any n, then from Theorem 1.7, (1.7) is obvious.

□

Example 1.8. For $r = 7, 11$, for all $n \geq 0$, $\alpha \geq 0$, we have

$$ped_{-2}\left(3^{2\alpha+2}n + \frac{r \times 3^{2\alpha+1} - 1}{4}\right) \equiv 0 \pmod{8}.$$

Combining Lin's Theorem 1.5, we have

$$ped_{-2}\left(3^{2\alpha+2}n + \frac{11 \times 3^{2\alpha+1} - 1}{4}\right) \equiv 0 \pmod{24}.$$

In section 2, we prove Theorem 1.2 and in section 3 we prove Theorem 1.7. Our method is elementary which is motivated by that of Mahlburg's [6]. We only use the knowledge of binary quadratic forms.

Acknowledgments.
We wish to thank the NSF of China (No.11071160) for its generous support. We would also like to thank the referee for his/her helpful comments.

2 Proof of Theorem 1.2

In this section, we prove Theorem 1.2. We need the following well-known identities [7]:

$$\prod_{n=1}^{\infty}\frac{(1-q^n)^2}{(1-q^{2n})} = \sum_{n=-\infty}^{\infty}(-1)^n q^{n^2} = 1 + 2\sum_{n=1}^{\infty}(-1)^n q^{n^2}, \tag{2.1}$$

$$q\prod_{n=1}^{\infty}\frac{(1-q^{16n})^2}{(1-q^{8n})} = \sum_{n=0}^{\infty} q^{(2n+1)^2}. \tag{2.2}$$

Proof of Theorem 1.2. Since

$$\sum_{n=0}^{\infty} ped(n)q^n := \prod_{n=1}^{\infty}\frac{(1-q^{4n})}{(1-q^n)} = \prod_{n=1}^{\infty}\frac{(1-q^{2n})^2}{(1-q^n)} \cdot \frac{(1-q^{4n})}{(1-q^{2n})^2},$$

so we have

$$\begin{aligned}
&\sum_{n=0}^{\infty} ped(n)q^{8n+1} \\
&= q\prod_{n=1}^{\infty}\frac{(1-q^{16n})^2}{1-q^{8n}}\cdot\frac{(1-q^{32n})}{(1-q^{16n})^2} \\
&= \left(\sum_{n=0}^{\infty} q^{(2n+1)^2}\right)\cdot\frac{1}{1+2\sum_{n=1}^{\infty}(-1)^n q^{16n^2}} \qquad (\text{by } (2.1),(2.2)) \\
&\equiv \left(\sum_{n=0}^{\infty} q^{(2n+1)^2}\right)\left(1-2\sum_{n=1}^{\infty}(-1)^n q^{16n^2}+4\left(\sum_{n=1}^{\infty}(-1)^n q^{16n^2}\right)^2\right) (\mathrm{mod}\, 8).
\end{aligned}$$

From above, it is clear that $ped(n)$ is odd if and only if $8n+1$ is a square. Note that

$$\begin{aligned}
\left(\sum_{n=1}^{\infty}(-1)^n q^{16n^2}\right)^2 &= \left(\sum_{m_1,m_2=1}^{\infty}(-1)^{m_1+m_2}q^{16m_1^2+16m_2^2}\right) \\
&= 2\left(\sum_{\substack{m_1,m_2=1\\ m_1<m_2}}^{\infty}(-1)^{m_1+m_2}q^{16m_1^2+16m_2^2}\right)+\left(\sum_{n=1}^{\infty}q^{32n^2}\right).
\end{aligned}$$

So we have

$$\begin{aligned}
\sum_{n=0}^{\infty} ped(n)q^{8n+1} &\equiv \left(1-2\sum_{n=1}^{\infty}(-1)^n q^{16n^2}+4\left(\sum_{n=1}^{\infty}q^{32n^2}\right)\right) \qquad (2.3)\\
&\quad\times\left(\sum_{n=0}^{\infty}q^{(2n+1)^2}\right) \quad (\mathrm{mod}\, 8). \qquad (2.4)
\end{aligned}$$

Note that if $8n+1 = p^{2\alpha+1}(8pm+r)$ where $p \equiv 7(\mathrm{mod}\, 8)$, $(r,p)=1$, $rp \equiv 1(\mathrm{mod}\, 8)$, then $8n+1$ can't be represented by x^2, x^2+y^2 or x^2+2y^2. So for these $8n+1$, from (2.3) and (2.4), it is easy to see that $ped(n) \equiv 0(\mathrm{mod}\, 8)$. This completes the proof of Theorem 1.2. □

3 Proof of Theorem 1.7

In this section, we prove Theorem 1.7. The method is similar to that of Theorem 1.2.

Proof of Theorem 1.7. Since

$$\sum_{n=0}^{\infty} ped_{-2}(n)q^n := \prod_{n=1}^{\infty} \frac{(1-q^{4n})^2}{(1-q^n)^2} = \prod_{n=1}^{\infty} \frac{(1-q^{4n})^2}{(1-q^{2n})} \cdot \frac{(1-q^{2n})}{(1-q^n)^2},$$

so by (2.1) and (2.2), we have

$$\begin{aligned}
\sum_{n=0}^{\infty} ped_{-2}(n)q^{4n+1} &= q\prod_{n=1}^{\infty} \frac{(1-q^{16n})^2}{1-q^{8n}} \cdot \frac{(1-q^{8n})}{(1-q^{4n})^2} \\
&= \frac{1}{1+2\sum_{n=1}^{\infty}(-1)^n q^{4n^2}} \times \left(\sum_{n=0}^{\infty} q^{(2n+1)^2}\right) \\
&\equiv \left(1-2\sum_{n=1}^{\infty}(-1)^n q^{4n^2} + 4\left(\sum_{n=1}^{\infty}(-1)^n q^{4n^2}\right)^2\right) \\
&\quad \times \left(\sum_{n=0}^{\infty} q^{(2n+1)^2}\right) \pmod 8.
\end{aligned}$$

Note that

$$\begin{aligned}
\left(\sum_{n=1}^{\infty}(-1)^n q^{4n^2}\right)^2 &= \left(\sum_{m_1,m_2=1}^{\infty}(-1)^{m_1+m_2} q^{4m_1^2+4m_2^2}\right) \\
&= 2\left(\sum_{\substack{m_1,m_2=1 \\ m_1<m_2}}^{\infty}(-1)^{m_1+m_2} q^{4m_1^2+4m_2^2}\right) + \sum_{n=1}^{\infty} q^{8n^2} \\
&= 2\left(\sum_{\substack{m_1,m_2=1 \\ m_1<m_2}}^{\infty}(-1)^{m_1+m_2} q^{4m_1^2+4m_2^2}\right) + \frac{1}{2}\left(\sum_{n=\infty}^{\infty} q^{8n^2} - 1\right).
\end{aligned}$$

Note also that

$$\begin{aligned}
\sum_{n=1}^{\infty}(-1)^n q^{4n^2} &= \frac{1}{2}\left(\sum_{n=-\infty}^{\infty}(-1)^n q^{4n^2} - 1\right) \\
&= -\frac{1}{2} - \frac{1}{2}\left(\sum_{n=-\infty}^{\infty} q^{4n^2}\right) + \left(\sum_{n=-\infty}^{\infty} q^{16n^2}\right).
\end{aligned}$$

So we have

$$\begin{aligned}\sum_{n=0}^{\infty} ped_{-2}(n)q^{4n+1} &\equiv \left(\sum_{n=-\infty}^{\infty} q^{4n^2} - 2\sum_{n=-\infty}^{\infty} q^{16n^2} + 2\sum_{n=\infty}^{\infty} q^{8n^2}\right) \\ &\quad \times\frac{1}{2}\left(\sum_{n=\infty}^{\infty} q^{(2n+1)^2}\right) \\ &\equiv \left(\frac{1}{2}\sum_{n=-\infty}^{\infty} q^{4n^2} + \sum_{n=\infty}^{\infty} q^{8n^2} - \sum_{n=-\infty}^{\infty} q^{16n^2}\right) \quad (3.1) \\ &\quad \times\left(\sum_{n=\infty}^{\infty} q^{(2n+1)^2}\right) \pmod 8. \quad (3.2)\end{aligned}$$

If we put

$$\left(\sum_{n=\infty}^{\infty} q^{(2n+1)^2}\right)\left(\sum_{n=\infty}^{\infty} q^{4n^2}\right) := \sum_{n=0}^{\infty} a(n)q^{4n+1}, \quad (3.3)$$

$$\left(\sum_{n=\infty}^{\infty} q^{(2n+1)^2}\right)\left(\sum_{n=\infty}^{\infty} q^{8n^2}\right) := \sum_{n=0}^{\infty} b(n)q^{4n+1}, \quad (3.4)$$

and

$$\left(\sum_{n=\infty}^{\infty} q^{(2n+1)^2}\right)\left(\sum_{n=\infty}^{\infty} q^{16n^2}\right) := \sum_{n=0}^{\infty} c(n)q^{4n+1}, \quad (3.5)$$

then it is easy to see from (3.1), (3.2) that

(1) $ped_{-2}(n)$ is odd if and only if $\frac{a(n)}{2}$ is odd,

(2) $ped_{-2}(n) \equiv 2 \pmod 4$ if and only if $\frac{a(n)}{2} + b(n) - c(n) \equiv 2 \pmod 4$,

(3) $ped_{-2}(n) \equiv 4 \pmod 8$ if and only if $\frac{a(n)}{2} + b(n) - c(n) \equiv 4 \pmod 8$,

(4) In all other cases, we have that $ped_{-2}(n) \equiv 0 \pmod 8$.

We separate into two cases according to the parity of n.
(i) If $n = 2m$ is even, then it is easy to see that

$$\begin{aligned}&\#\{(x,y) \in \mathbb{Z}\times\mathbb{Z} : 8m+1 = x^2 + 16y^2, 8m+1 \text{ is not a square}\} \\ =\ &\#\{(x,y) \in \mathbb{Z}\times\mathbb{Z} : 8m+1 = x^2 + 4y^2, 8m+1 \text{ is not a square}\} \\ =\ &\frac{1}{2}\#\{(x,y) \in \mathbb{Z}\times\mathbb{Z} : 8m+1 = x^2 + y^2, 8m+1 \text{ is not a square}\},\end{aligned}$$

and

$$\begin{aligned} & \#\{(x,y)\in\mathbb{Z}\times\mathbb{Z} : 8m+1 = x^2+8y^2, 8m+1 \text{ is not a square}\} \\ = \; & \#\{(x,y)\in\mathbb{Z}\times\mathbb{Z} : 8m+1 = x^2+2y^2, 8m+1 \text{ is not a square}\} \end{aligned}$$

Note that if $8m+1 = p_1^{\alpha_1}\cdots p_k^{\alpha_k} q_1^{\beta_1}\cdots q_l^{\beta_l} r_1^{\gamma_1}\cdots r_u^{\gamma_u} s_1^{\delta_1}\cdots s_v^{\delta_v}$, where $p_i \equiv 1 \pmod 8$, $q_i \equiv 3 \pmod 8$, $r_i \equiv 5 \pmod 8$ and $s_i \equiv 7 \pmod 8$, then by using the decomposition of prime ideals in $\mathbb{Z}[i]$, we know that $8m+1 = x^2+y^2$ has integral solutions if and only if $\beta_j, \gamma_t \equiv 0 \pmod 2$ for $1 \le j \le l$ and $1 \le t \le u$. If all β_j, δ_t are even, then it is easy to see that

$$\#\{(x,y)\in\mathbb{Z}\times\mathbb{Z} : 8m+1 = x^2+y^2\} = 4(\alpha_1+1)\cdots(\alpha_k+1)(\gamma_1+1)\cdots(\gamma_u+1).$$

We obtain that $a(n) = 2(\alpha_1+1)\cdots(\alpha_k+1)(\gamma_1+1)\cdots(\gamma_u+1)$ if all β_j, γ_t are even. Now it is clear that $\frac{a(n)}{2}$ is odd if and only if $4n+1 = 8m+1$ is a square.

Similarly, by using the decomposition of prime ideals in $\mathbb{Z}[\sqrt{-2}]$, we know that if all $\gamma_j, \delta_t \equiv 0 \pmod 2$, then (there are only two roots of unities ± 1 in $\mathbb{Z}[\sqrt{-2}]$)

$$\#\{(x,y)\in\mathbb{Z}\times\mathbb{Z} : 8m+1 = x^2+2y^2\} = 2(\alpha_1+1)\cdots(\alpha_k+1)(\beta_1+1)\cdots(\beta_l+1).$$

From the above argument, we obtain the following results. Suppose all δ_i are even, then

(a) If β_i is odd for some $1 \le i \le l$ and all γ_j is even, then $\frac{a(n)}{2} + b(n) - c(n) \equiv b(n) \equiv 2(\alpha_1+1)\cdots(\alpha_k+1)(\beta_1+1)\cdots(\beta_l+1) \equiv 0 \pmod 8$ (because there must be another $\beta_{i'}$ odd for $p_i \equiv 3 \pmod 8$ while $8m+1 \equiv 1 \pmod 8$).

(b) If γ_i is odd for some $1 \le i \le u$ and all β_j is even, then $\frac{a(n)}{2} + b(n) - c(n) \equiv \frac{a(n)}{2} - c(n) \equiv -(\alpha_1+1)\cdots(\alpha_k+1)(\gamma_1+1)\cdots(\gamma_u+1) \pmod 8$.

(c) If all β_i and γ_j are even, then $\frac{a(n)}{2} + b(n) - c(n) \equiv -(\alpha_1+1)\cdots(\alpha_k+1)(\gamma_1+1)\cdots(\gamma_u+1) + 2(\alpha_1+1)\cdots(\alpha_k+1)(\beta_1+1)\cdots(\beta_l+1) \pmod 8$.

(d) If β_i and γ_j are odd for some i and j, then clearly $ped_{-2}(n) \equiv 0 \pmod 8$.

From (a),(b),(c) and (d), it is not difficult to see that (for $n = 2m$)

(1) if $4n+1$ is a square, then $ped_{-2}(n)$ is odd,

(2) if $4n+1 = pa^2$ where $\text{ord}_p(4n+1) \equiv 1 (\text{mod}\, 4)$, then $ped_{-2}(n) \equiv 2 (\text{mod}\, 4)$,

(3) if $4n+1 = p_1 p_2 a^2$ where $p_i \equiv 1 (\text{mod}\, 4)$ and $\text{ord}_{p_i}(4n+1) \equiv 1 (\text{mod}\, 4)$ or $4n+1 = p^3 a^2$ and $\text{ord}_p(4n+1) \equiv 3 (\text{mod}\, 8)$, then $ped_{-2}(n) \equiv 4 (\text{mod}\, 8)$

(4) and for all other cases, $ped_{-2}(n) \equiv 0 (\text{mod}\, 8)$.

(ii) If $n = 2m+1$, then $4n+1 = 8m+5$. Since $(2m_1+1)^2 + 8m_2^2 \equiv (2m_1+1)^2 + 16m_2^2 \equiv 1 (\text{mod}\, 8)$, so $4n+1$ can not be represented by $(2m_1+1)^2 + 8m_2^2$ and $(2m_1+1)^2 + 16m_2^2$. Note also that

$$\begin{aligned} & \#\{(x,y) \in \mathbb{Z} \times \mathbb{Z} : 8m+5 = x^2 + 4y^2\} \\ = \ & \frac{1}{2} \#\{(x,y) \in \mathbb{Z} \times \mathbb{Z} : 8m+5 = x^2 + y^2\}. \end{aligned}$$

So if $8m+5 = p_1^{\alpha_1} \cdots p_k^{\alpha_k} q_1^{\beta_1} \cdots q_l^{\beta_l} r_1^{\gamma_1} \cdots r_u^{\gamma_u} s_1^{\delta_1} \cdots s_v^{\delta_v}$, where $p_i \equiv 1 (\text{mod}\, 8)$, $q_i \equiv 3 (\text{mod}\, 8)$, $r_i \equiv 5 (\text{mod}\, 8)$ and $s_i \equiv 7 (\text{mod}\, 8)$, then

$$\frac{a(n)}{2} \equiv (\alpha_1 + 1) \cdots (\alpha_k + 1)(\gamma_1 + 1) \cdots (\gamma_u + 1) \pmod 8. \tag{3.6}$$

Just like in (i), it is easy to find case by case by (3.6) for $ped_n (\text{mod}\, 8)$ for $n = 2m+1$:

(1) if $4n+1 = pa^2$ where $\text{ord}_p(4n+1) \equiv 1 (\text{mod}\, 4)$, then $ped_{-2}(n) \equiv 2 (\text{mod}\, 4)$,

(2) if $4n+1 = p_1 p_2 a^2$ and $\text{ord}_{p_i}(4n+1) \equiv 1 (\text{mod}\, 4)$ or $4n+1 = p^3 a^2$ and $\text{ord}_p(4n+1) \equiv 3 (\text{mod}\, 8)$, then $ped_{-2}(n) \equiv 4 (\text{mod}\, 8)$

(3) and for all other cases, $ped_{-2}(n) \equiv 0 (\text{mod}\, 8)$.

Now combining (i) and (ii), with a little thought, we complete the proof of Theorem 1.7.

□

References

[1] G .E. Andrews, M. D. Hirschhorn and J.A. Sellers *Arithmetic properties of partitions with even parts distinct*. Ramanujan J. **22** (2010), pp. 273-284.

[2] A. Berkovich and H. Yesilyurt; *Ramanujan's identities and representation of integers by certain binary and quaternary quadratic forms*. Ramanujan J. **20** (2009), no. 3, 375-408.

[3] S. C. Chen, *On the number of partitions with distinct even parts*. Discrete Math. **311** (2011), no. 12, 940-943.

[4] S.C.Chen, *Congruences for the number of k-tuple partitions with distinct even parts*, Discrete Math. **313** (2013), 1565-1568.

[5] Bernard L. S. Lin, *Arithmetic properties of bipartitions with even parts distinct*, Ramanujan J. To appear.

[6] K. Mahlburg, *The overpartition function modulo small powers of 2*, Discrete Math. **286** (2004), no. 3, 263-267.

[7] K. Ono, *The web of modularity: arithmetic of the coefficients of modular forms and q-series*, CBMS Regional Conference Series in Mathematics, **102** American Mathematical Society, Providence, RI, 2004.

[8] K. Ono and D. Penniston, *The 2-adic behavior of the number of partitions into distinct parts*, J. Combin. Theory (Ser.A) **92** (2000), 138-157.